Amendments to the Claims

Listing of Claims:

Claim 1 (original): A method of treating mesothelioma comprising administering a therapeutically effective amount of a 4-pyridylmethyl-phthalazine derivative to a warmblooded animal in need thereof.

Claim 2 (original): Method according to claim 1 comprising administering a therapeutically effective amount of a 4-pyridylmethyl-phthalazine derivative of formula I

wherein

r is 0 to 2,

n is 0 to 2,

m is 0 to 4,

R₁ and R₂ (i) are lower alkyl or

(ii) together form a bridge in subformula I*

$$(I^*)$$

the binding being achieved via the two terminal carbon atoms, or

(iii) together form a bridge in subformula !**

$$\begin{array}{ccc}
T_1 \\
T_2 \\
T_4 = T_3
\end{array} \tag{I**}$$

wherein one or two of the ring members T_1 , T_2 , T_3 and T_4 are nitrogen, and the others are in each case CH, and the binding is achieved via T_1 and T_4 ;

A, B, D, and E are, independently of one another, N or CH, with the stipulation that not

more than 2 of these radicals are N;

G is lower alkylene, lower alkylene substituted by acyloxy or hydroxy, -CH₂-O-, -CH₂-S-, -CH₂-NH-, oxa (-O-), thia (-S-), or imino (-NH-);

Q is lower alkyl;

R is H or lower alkyl;

X is imino, oxa, or thia;

Y is unsubstituted or substituted aryl, pyridyl, or unsubstituted or substituted cycloalkyl; and

Z is amino, mono- or disubstituted amino, halogen, alkyl, substituted alkyl, hydroxy, etherified or esterified hydroxy, nitro, cyano, carboxy, esterified carboxy, alkanoyl, carbamoyl, N-mono- or N,N-disubstituted carbamoyl, amidino, guanidino, mercapto, sulfo, phenylthio, phenyl-lower alkylthio, alkylphenylthio, phenylsulfonyl, phenyl-lower alkylsulfinyl or alkylphenylsulfinyl, substituents Z being the same or different from one another if more than 1 radical Z is present;

and wherein the bonds characterized, if present, by a wavy line are either single or double bonds:

or an N-oxide of the defined compound, wherein 1 or more N atoms carry an oxygen atom, or the salt of such compound having at least one salt-forming group, to a warm-blooded animal in need thereof.

Claim 3 (original): Method of claim 2 wherein the 4-pyridylmethyl-phthalazine derivative of formula I is 1-(4-chloroanilino)-4-(4-pyridylmethyl)phthalazine.

Claim 4 (currently amended): Method according to any one of claims 1 to 3 claim 1 wherein the disease is selected from pleural mesothelioma, peritoneal mesothelioma, pericardial mesothelioma, epithelial mesothelioma, sarcomatous mesothelioma and biphasic mesothelioma.

Claim 5 (currently amended): Method according to any one of claims 1 to 4 claim 1 wherein the warm-blooded animal is a human.

Claim 6 (original): Method according to claim 5 which comprises administering 1-(4-chloroanilino)-4-(4-pyridylmethyl)phthalazine, or a pharmaceutically acceptable salt thereof, to the patient on a once daily schedule at a dose in the range from 1000 mg/day to 1400 mg/day.

Claim 7 (original): Method according to claim 6 wherein the once daily dose is 1200 mg/day to 1300 mg/day.

Claim 8 (original): Method according to claim 6 wherein the once daily dose is 1250 mg/day.

Claim 9 (original): A method of treating mesothelioma comprising administering a 4-pyridylmethyl-phthalazine derivative in an amount which is therapeutically effective against mesothelioma to a warm-blooded animal in need thereof in combination with surgery, in particular pleurectomy/decortication, pneumonectomy, extrapleural pneumonectomy, and/or radiation therapy, in particular external radiation therapy or internal radiation therapy,

Claim 10 (original): A commercial package comprising a 4-pyridylmethyl-phthalazine derivative together with instructions for use thereof in the treatment of mesothelioma.

Claim 11 (canceled)

Claim 12 (currently amended): <u>Use-Method</u> according to claim <u>442</u>, wherein the 4-pyridylmethyl-phthalazine derivative is 1-(4-chloroanilino)-4-(4-pyridylmethyl)phthalazine or 1-(4-chloroanilino)-4-(4-pyridylmethyl)phthalazine in free form or in the form of a pharmaceutically acceptable salt.

Claim 13 (currently amended): <u>Use-Method</u> according to claim 12, wherein the pharmaceutically acceptable salt is the succinate salt of the 4-pyridylmethyl-phthalazine derivative.